
| | |
|--|---------|
| <i>Ageniella fabricii</i> Banks, 1946 | 1 ex. |
| <i>Agenioideus apicalis</i> Van der Linden, 1827 | 1 ex. |
| <i>Agenioideus ciliatus</i> Lepeletier, 1845 | 1 ex. |
| <i>Agenioideus cinctellus</i> Spinola, 1808 | 169 ex. |
| <i>Agenioideus dichrous</i> Brullé, 1840 | 1 ex. |
| <i>Agenioideus excisus</i> Morawitz, 1890 | 1 ex. |
| <i>Agenioideus humilis</i> Cresson, 1867 | 1 ex. |
| <i>Agenioideus nubecula</i> Costa, 1881 | 6 ex. |
| <i>Agenioideus sericeus</i> Van der Linden, 1827 | 63 ex. |
| <i>Agenioideus spec.</i> | 3 ex. |
| <i>Agenioideus usurarius</i> Tournier, 1889 | 4 ex. |
| <i>Allaporus pulchellus</i> Banks, 1910 | 1 ex. |
| <i>Allocyphonyx maura</i> Cresson | 1 ex. |
| <i>Anoplius aeruginosus</i> Tournier, 1890 | 2 ex. |
| <i>Anoplius aethiops</i> Cresson, 1865 | 14 ex. |
| <i>Anoplius amethystinus</i> Fabricius, 1793 | 9 ex. |
| <i>Anoplius apiculatus pretiosus</i> Banks, 1910 | 1 ex. |
| <i>Anoplius araucanus</i> Herbst | 1 ex. |
| <i>Anoplius atrox</i> Dahlbom | 1 ex. |
| <i>Anoplius beatus</i> Cameron | 1 ex. |
| <i>Anoplius canifrons</i> Smith | 11 ex. |
| <i>Anoplius caviventris</i> Aurivillius, 1907 | 6 ex. |
| <i>Anoplius clystera</i> Banks, 1914 | 10 ex. |
| <i>Anoplius compactus</i> Banks | 4 ex. |
| <i>Anoplius concinnus</i> Dahlbom, 1845 | 32 ex. |
| <i>Anoplius cylindricus</i> Cresson, 1867 | 2 ex. |
| <i>Anoplius diphonichus</i> Spinola | 15 ex. |
| <i>Anoplius dispar</i> Dahlbom | 26 ex. |
| <i>Anoplius dumerosus</i> Spinola | 8 ex. |
| <i>Anoplius eureka</i> Banks | 4 ex. |
| <i>Anoplius fazii</i> Herbst, i.l. | 2 ex. |
| <i>Anoplius fragilis</i> Smith | 3 ex. |
| <i>Anoplius ignitus</i> Smith | 1 ex. |
| <i>Anoplius ilione</i> Banks | 2 ex. |
| <i>Anoplius illinoiensis</i> Rbt. | 2 ex. |
| <i>Anoplius imbellis</i> Banks, 1944 | 2 ex. |
| <i>Anoplius infuscatus</i> Van der Linden, 1827 | 125 ex. |
| <i>Anoplius insidiosus</i> Smith | 5 ex. |
| <i>Anoplius iwatai</i> Yasumatsu | 2 ex. |
| <i>Anoplius linkin</i> Dalla Torre | 3 ex. |
| <i>Anoplius marginalis</i> Banks, 1910 | 1 ex. |
| <i>Anoplius marginatus</i> Say, 1824 | 11 ex. |
| <i>Anoplius misturatus</i> Kohl | 12 ex. |
| <i>Anoplius morosus</i> Smith | 2 ex. |
| <i>Anoplius nigerrimus</i> Scopoli, 1763 | 105 ex. |
| <i>Anoplius nigritus</i> Dahlbom, 1843 | 2 ex. |
| <i>Anoplius nitidulus</i> Guérin | 13 ex. |

| | |
|--|---------|
| <i>Anoplius perplexa</i> Smith | 2 ex. |
| <i>Anoplius petiolaris</i> Gussakovskij, 1929 | 1 ex. |
| <i>Anoplius phalerata</i> Perty | 1 ex. |
| <i>Anoplius puniceus</i> Sustera, i.l. | 2 ex. |
| <i>Anoplius reflexus</i> Smith | 1 ex. |
| <i>Anoplius samariensis</i> Pallas, 1771 | 4 ex. |
| <i>Anoplius schlettereri</i> Radoszkowsky, 1888 | 2 ex. |
| <i>Anoplius semirufus</i> Haupt | 1 ex. |
| <i>Anoplius separatus</i> Taschenberg | 2 ex. |
| <i>Anoplius smaragdinus</i> Herbst | 2 ex. |
| <i>Anoplius sobrinus</i> Spinola | 8 ex. |
| <i>Anoplius solanus</i> Kohl | 3 ex. |
| <i>Anoplius</i> sp. | 2 ex. |
| <i>Anoplius tenebrosus</i> Cresson, 1865 | 1 ex. |
| <i>Anoplius tenuicornis</i> Tournier, 1889 | 4 ex. |
| <i>Anoplius thoracicus</i> Rossi | 2 ex. |
| <i>Anoplius ventralis</i> Banks, 1910 | 1 ex. |
| <i>Anoplius veranes</i> Banks | 2 ex. |
| <i>Anoplius vespiformis</i> Klug | 9 ex. |
| <i>Anoplius viaticus</i> Linné, 1758 | 347 ex. |
| <i>Anoplius viaticus haupti</i> Guiglia | 7 ex. |
| <i>Anoplius viaticus paganus</i> Dahlbom, 1843 | 21 ex. |
| <i>Anoplius virginiensis</i> Cresson | 2 ex. |
| <i>Anospilus ichnusus</i> Wolf, 1962 | 2 ex. |
| <i>Anospilus melanarius</i> Van der Linden, 1827 | 2 ex. |
| <i>Anospilus orbitalis luctiger</i> Costa, 1886 | 5 ex. |
| <i>Anospilus orbitalis orbitalis</i> Costa, 1863 | 13 ex. |
| <i>Anospilus sardus</i> Priesner, 1962 | 1 ex. |
| <i>Aporinellus apicatus</i> Banks | 2 ex. |
| <i>Aporinellus apicipennis</i> Bréthes | 1 ex. |
| <i>Aporinellus hecate</i> Cameron, 1891 | 1 ex. |
| <i>Aporinellus sexmaculatus</i> Spinola, 1805 | 12 ex. |
| <i>Aporus bicolor</i> Spinola, 1808 | 8 ex. |
| <i>Aporus bicolor fulviventris</i> Costa, 1882 | 1 ex. |
| <i>Aporus luxus</i> Banks, 1914 | 6 ex. |
| <i>Aporus luxus assimilis</i> Banks, 1917 | 1 ex. |
| <i>Aporus planiceps</i> Latreille, 1809 | 1 ex. |
| <i>Aporus pollux</i> Kohl, 1888 | 16 ex. |
| <i>Aporus unicolor</i> Spinola, 1808 | 27 ex. |
| <i>Arachnospila abnormis</i> Dahlbom, 1842 | 18 ex. |
| <i>Arachnospila alvarabnormis</i> Wolf, 1965 | 1 ex. |
| <i>Arachnospila anceps</i> Wesmael, 1851 | 398 ex. |
| <i>Arachnospila angularis</i> Banks, 1910 | 1 ex. |
| <i>Arachnospila ausa</i> Tournier, 1890 | 5 ex. |
| <i>Arachnospila consobrina</i> Dahlbom, 1843 | 35 ex. |
| <i>Arachnospila esau</i> Kohl, 1886 | 2 ex. |
| <i>Arachnospila fumipennis</i> Zetterstedt, 1838 | 10 ex. |

| | |
|---|---------|
| <i>Arachnospila fumipennis paladeae</i> Wolf & Moczar, 1972 | 5 ex. |
| <i>Arachnospila fuscomarginata</i> Thomson, 1870 | 34 ex. |
| <i>Arachnospila hedickei</i> Haupt, 1929 | 1 ex. |
| <i>Arachnospila minutula</i> Dahlbom, 1842 | 70 ex. |
| <i>Arachnospila proximus mongoloproximus</i> Wolf & Mocsáry | 1 ex. |
| <i>Arachnospila pseudabnormis</i> Wolf, 1965 | 11 ex. |
| <i>Arachnospila rhaetabnormis</i> Wolf, 1965 | 1 ex. |
| <i>Arachnospila rufa</i> Haupt, 1927 | 22 ex. |
| <i>Arachnospila sogdiana</i> Morawitz, 1893 | 16 ex. |
| <i>Arachnospila</i> sp. | 1 ex. |
| <i>Arachnospila spissa</i> Schiödte, 1837 | 291 ex. |
| <i>Arachnospila trivialis</i> Dahlbom, 1843 | 253 ex. |
| <i>Arachnospila usurata</i> Blüthgen, 1957 | 15 ex. |
| <i>Arachnospila vasumatsui</i> Wolf & Mocsáry | 1 ex. |
| <i>Arachnospila virgilabnormis</i> Wolf, 1976 | 2 ex. |
| <i>Arachnospila wesmaeli</i> Thomson, 1870 | 21 ex. |
| <i>Arachnospila westerlundi</i> Morawitz, 1893 | 1 ex. |
| <i>Auplopus aegina</i> Smith, 1857 | 16 ex. |
| <i>Auplopus alaris</i> Saussure, 1867 | 8 ex. |
| <i>Auplopus albifrons</i> Dalman, 1823 | 14 ex. |
| <i>Auplopus architectus metallicus</i> Banks, 1910 | 4 ex. |
| <i>Auplopus blanda</i> Guérin | 2 ex. |
| <i>Auplopus caerulea</i> Smith | 3 ex. |
| <i>Auplopus callisto</i> Smith, 1859 | 2 ex. |
| <i>Auplopus canaliculata</i> Cameron | ex. |
| <i>Auplopus carbonarius</i> Scopoli, 1763 | 232 ex. |
| <i>Auplopus celaeno</i> Smith, 1857 | 1 ex. |
| <i>Auplopus chrysosoma</i> Rohwer, 1910 | 1 ex. |
| <i>Auplopus cinnamomea</i> Arnold, 1936 | 2 ex. |
| <i>Auplopus clypeata</i> Bingham, 1896 | 5 ex. |
| <i>Auplopus commendabilis</i> Kohl, 1894 | 3 ex. |
| <i>Auplopus deceptrix</i> Smith | 7 ex. |
| <i>Auplopus ferruginea</i> Najo | 2 ex. |
| <i>Auplopus flavotegulatus</i> Bingham, 1902 | 1 ex. |
| <i>Auplopus formosana</i> Susterka, i.l. | 4 ex. |
| <i>Auplopus hessei</i> Arnold | 1 ex. |
| <i>Auplopus jacobi</i> Arnold | 1 ex. |
| <i>Auplopus kankau</i> Babiy, i.l. | 21 ex. |
| <i>Auplopus laeviculus</i> Bingham, 1897 | 2 ex. |
| <i>Auplopus mellipes</i> Say, 1836 | 1 ex. |
| <i>Auplopus mygnimioides</i> Bisch. | 3 ex. |
| <i>Auplopus nebulosus</i> Dreisbach | 2 ex. |
| <i>Auplopus nobilis</i> Fabricius | 1 ex. |
| <i>Auplopus novarae</i> Saussure | 28 ex. |
| <i>Auplopus personata</i> Gribodo, 1879 | 2 ex. |
| <i>Auplopus personata ornaticollis</i> Cameron | 2 ex. |
| <i>Auplopus pseudocallisto</i> Babiy, i.l. | 3 ex. |

| | |
|---|---------|
| <i>Auplopus pseudotincta</i> Suster, i.l. | 1 ex. |
| <i>Auplopus pulchricoma</i> Arnold, 1934 | 1 ex. |
| <i>Auplopus pygialis</i> Pérez, 1905 | 1 ex. |
| <i>Auplopus rectus</i> Haupt, 1927 | 8 ex. |
| <i>Auplopus rubronigra</i> Suster, i.l. | 1 ex. |
| <i>Auplopus sauteri</i> Suster, i.l. | 1 ex. |
| <i>Auplopus</i> sp. | 1 ex. |
| <i>Auplopus spilotoenia</i> Kohl, 1894 | 1 ex. |
| <i>Auplopus suster</i> Babiy, i.l. | 1 ex. |
| <i>Auplopus taihorinensis</i> Babiy, i.l. | 1 ex. |
| <i>Auplopus takaehioi</i> Yasumatsu | 1 ex. |
| <i>Auplopus tincta</i> Smith | 15 ex. |
| <i>Auplopus vaga pallidipennis</i> Arnold | 1 ex. |
| <i>Auplopus varitarsata</i> Dalla Torre | 1 ex. |
| <i>Batozonus unifasciatus</i> Smith | 1 ex. |
| <i>Batozonellus annulatus</i> Fabricius, 1793 | 5 ex. |
| <i>Batozonellus capensis</i> Dahlbom, 1843 | 4 ex. |
| <i>Batozonellus fuliginosus</i> Klug, 1834 | 6 ex. |
| <i>Batozonellus lacerticida</i> Pallas, 1771 | 72 ex. |
| <i>Batozonellus maculifrons</i> Smith, 1873 | 4 ex. |
| <i>Batozonus pygidialis</i> Kohl | 1 ex. |
| <i>Caliadurgus fasciatellus</i> Spinola, 1808 | 320 ex. |
| <i>Ceropales albicincta</i> Rossi, 1790 | 7 ex. |
| <i>Ceropales albicincta seraxensis</i> Radoszkowsky, 1893 | 2 ex. |
| <i>Ceropales altaica</i> Morawitz, 1888 | 1 ex. |
| <i>Ceropales bipunctata</i> Say, 1824 | 1 ex. |
| <i>Ceropales chilensis</i> Spinola, 1851 | 26 ex. |
| <i>Ceropales fraterna</i> Say | 10 ex. |
| <i>Ceropales judicatrix</i> Nurse, 1902 | 1 ex. |
| <i>Ceropales maculata</i> Fabricius, 1775 | 117 ex. |
| <i>Ceropales picta</i> Shuckard, 1837 | 1 ex. |
| <i>Ceropales pygmaea</i> Kohl, 1880 | 1 ex. |
| <i>Ceropales</i> sp. | 1 ex. |
| <i>Ceropales variegata</i> Fabricius, 1798 | 20 ex. |
| <i>Chalcochaes hirsutifemur</i> Banks, 1914 | 2 ex. |
| <i>Chirodamus pachylopus</i> Kohl | 2 ex. |
| <i>Clavelia bicolor</i> Smith | 1 ex. |
| <i>Cryptocheilus alternatus</i> Lepeletier, 1845 | 3 ex. |
| <i>Cryptocheilus australis</i> Guérin | 11 ex. |
| <i>Cryptocheilus brunneus</i> Klug | 2 ex. |
| <i>Cryptocheilus coloratus</i> Suster, 1924 | 0 ex. |
| <i>Cryptocheilus discolor</i> Fabricius, 1793 | 5 ex. |
| <i>Cryptocheilus egregius</i> Lepeletier, 1845 | 4 ex. |
| <i>Cryptocheilus elegans</i> Spinola, 1806 | 1 ex. |
| <i>Cryptocheilus elegans tripunctatus</i> Spinola | 0 ex. |
| <i>Cryptocheilus fabricii</i> Van der Linden, 1827 | 64 ex. |
| <i>Cryptocheilus fabricii manchurianus</i> Yasumatsu | 1 ex. |

| | |
|---|---------|
| <i>Cryptocheilus fischeri</i> Spinola | 2 ex. |
| <i>Cryptocheilus flammipennis</i> Smith | 1 ex. |
| <i>Cryptocheilus freygessneri</i> Kohl | 5 ex. |
| <i>Cryptocheilus fulvicollis</i> Costa, 1874 | 0 ex. |
| <i>Cryptocheilus guttulatus</i> Costa, 1886 | 7 ex. |
| <i>Cryptocheilus ichneumonoides</i> Costa, 1874 | 11 ex. |
| <i>Cryptocheilus infumatus</i> Palma, 1869 | 1 ex. |
| <i>Cryptocheilus lividocincta</i> Costa? | 0 ex. |
| <i>Cryptocheilus notatus</i> Rossi, 1792 | 35 ex. |
| <i>Cryptocheilus notatus affinis</i> Van der Linden, 1827 | 36 ex. |
| <i>Cryptocheilus octomaculatus</i> Rossi, 1790 | 3 ex. |
| <i>Cryptocheilus pseudonotatus</i> Susterera | 1 ex. |
| <i>Cryptocheilus richardsi</i> Moczar, 1953 | 2 ex. |
| <i>Cryptocheilus rogenhoferi</i> Radovskovsky | 1 ex. |
| <i>Cryptocheilus rubellus</i> Eversmann, 1846 | 6 ex. |
| <i>Cryptocheilus semicastaneus</i> Morawitz, 1894 | 3 ex. |
| <i>Cryptocheilus</i> sp. | 1 ex. |
| <i>Cryptocheilus</i> spec. | 3 ex. |
| <i>Cryptocheilus terminatum</i> Say, 1828 | 1 ex. |
| <i>Cryptocheilus usambarensis</i> Lucas | 2 ex. |
| <i>Cryptocheilus variabilis</i> Rossi, 1790 | 12 ex. |
| <i>Cryptocheilus variegatus</i> Fabricius, 1776 | 0 ex. |
| <i>Cryptocheilus versicolor</i> Scopoli, 1763 | 102 ex. |
| <i>Ctenegenia vespiformis</i> Klug | 3 ex. |
| <i>Cyphononyx aeneipennis</i> Lucas, 1898 | 1 ex. |
| <i>Cyphononyx aspasius</i> Smith | 1 ex. |
| <i>Cyphononyx auropubens</i> Arnold | 1 ex. |
| <i>Cyphononyx dorsalis</i> Lepeletier | 1 ex. |
| <i>Cyphononyx flavicornis antennatus</i> Smith, 1855 | 10 ex. |
| <i>Cyphononyx fulvognathus</i> Rohwer | 8 ex. |
| <i>Cyphononyx iridipennis</i> Smith | 1 ex. |
| <i>Cyphononyx optimus</i> Smith, 1855 | 1 ex. |
| <i>Cyphononyx splendens</i> Lucas | 1 ex. |
| <i>Dentagenia formosana</i> Wahis | 1 ex. |
| <i>Deuteragenia formosana</i> Susterera, i.l. | 1 ex. |
| <i>Deuteragenia orientalis</i> Susterera, i.l. | 4 ex. |
| <i>Dicyrtomellus argenteus</i> Wahis, 1982 | 5 ex. |
| <i>Dicyrtomellus</i> sp. | 1 ex. |
| <i>Dicyrtomus leptacantius</i> Cameron | 1 ex. |
| <i>Dipogon bifasciatus</i> Geoffroy, 1785 | 43 ex. |
| <i>Dipogon calipterus</i> Say, 1836 | 2 ex. |
| <i>Dipogon hyalinatus</i> Dahlbom | 3 ex. |
| <i>Dipogon hyrcanus</i> Fabricius | 1 ex. |
| <i>Dipogon monticolus</i> Wahis, 1972 | 4 ex. |
| <i>Dipogon nipponica</i> Yasumatsu | 1 ex. |
| <i>Dipogon nitidus</i> Haupt | 1 ex. |
| <i>Dipogon subintermedius</i> Magretti, 1886 | 238 ex. |

| | |
|--|---------|
| Dipogon variegatus Linné, 1758 | 8 ex. |
| Dipogon vechti Day, 1979 | 2 ex. |
| Eidopompilus kruegeri Kohl | 1 ex. |
| Entomobora affinis Haupt | 1 ex. |
| Entomobora bucephala Tournier | 5 ex. |
| Entomobora crassitarsis Costa, 1874 | 5 ex. |
| Entomobora crassitarsis damryi Tournier, 1889 | 1 ex. |
| Entomobora crassitarsis punica Haupt, 1937 | 1 ex. |
| Entomobora plicata Costa, 1883 | 2 ex. |
| Entomobora vomeriventrtris Costa, 1881 | 1 ex. |
| Entypus bituberculatus Guérin | 1 ex. |
| Entypus caeruleus Linné, 1758 | 1 ex. |
| Entypus carinatus Fox | 1 ex. |
| Entypus fulvicornis Cresson, 1867 | 2 ex. |
| Entypus taschenbergi Dalla Torre, 1895 | 9 ex. |
| Entypus unifasciatus Say, 1828 | 1 ex. |
| Entypus unifasciatus californicus Townes, 1957 | 2 ex. |
| Entypus unifasciatus dumosus Spinola | 17 ex. |
| Entypus unifasciatus ferrugineipennis Haliday | 1 ex. |
| Eoferreola manticata Pallas, 1771 | 1 ex. |
| Eoferreola rhombica Christ, 1791 | 32 ex. |
| Epipompilus sp. | 1 ex. |
| Episyron albonotatum Van der Linden, 1827 | 158 ex. |
| Episyron arrogans Smith, 1873 | 13 ex. |
| Episyron biguttatus Fabricius, 1798 | 11 ex. |
| Episyron conterminus Smith | 1 ex. |
| Episyron gallicum intermedium Haupt, 1930 | 4 ex. |
| Episyron gryps Saussure, 1892 | 3 ex. |
| Episyron orbitalis Haupt, 1962 | 1 ex. |
| Episyron quadrispinosum Wahis | 5 ex. |
| Episyron quinquenotatus Say, 1835 | 5 ex. |
| Episyron rufipes Linné, 1758 | 270 ex. |
| Episyron spec. | 2 ex. |
| Episyron suawi Viereck | 1 ex. |
| Episyron vagabundus Smith, 1857 | 2 ex. |
| Euclavelia fasciata Smith | 1 ex. |
| Euplaniceps saussurei Kohl | 1 ex. |
| Evagetes alamannicus Blüthgen, 1944 | 6 ex. |
| Evagetes crassicornis Shuckard, 1835 | 211 ex. |
| Evagetes dubius Van der Linden, 1827 | 79 ex. |
| Evagetes elongatus Lepeletier, 1845 | 1 ex. |
| Evagetes gibbulus Lepeletier, 1845 | 11 ex. |
| Evagetes iconionus Wolf, 1970 | 1 ex. |
| Evagetes littoralis Wesmael, 1851 | 4 ex. |
| Evagetes parvus Cresson, 1865 | 2 ex. |
| Evagetes pectinipes Linné, 1758 | 59 ex. |
| Evagetes proximus Dahlbom, 1843 | 23 ex. |

| | |
|--|--------|
| <i>Evagetes sahlbergi</i> Morawitz, 1893 | 8 ex. |
| <i>Evagetes siculus</i> Lepeletier, 1845 | 23 ex. |
| <i>Evagetes</i> sp. | 1 ex. |
| <i>Evagetes subangulatus</i> Banks, 1919 | 9 ex. |
| <i>Evagetes subglaber</i> Haupt, 1941 | 1 ex. |
| <i>Evagetes tumidosus</i> Tournier, 1890 | 4 ex. |
| <i>Evagetes tumidosus sinuatus</i> Haupt, 1941 | 3 ex. |
| <i>Ferreola diffinis</i> Lepeletier, 1845 | 3 ex. |
| <i>Ferreola fulvicollis</i> Guérin, 1844 | 2 ex. |
| <i>Ferreola sagax</i> Kohl, 1886 | 3 ex. |
| <i>Hemipepsis ascensi</i> Zavattari, 1907 | 1 ex. |
| <i>Hemipepsis barbara</i> Lepeletier, 1845 | 14 ex. |
| <i>Hemipepsis brunneiceps pringleae</i> Cameron | 1 ex. |
| <i>Hemipepsis caelebs</i> Arnold | 7 ex. |
| <i>Hemipepsis caelebs flavida</i> Arnold | 8 ex. |
| <i>Hemipepsis capensis</i> Fabricius | 2 ex. |
| <i>Hemipepsis convexa</i> Bingham | 1 ex. |
| <i>Hemipepsis dedjas</i> Guérin | 1 ex. |
| <i>Hemipepsis dimidiata</i> Wahis | 2 ex. |
| <i>Hemipepsis ducalis</i> Smith | 1 ex. |
| <i>Hemipepsis fallax</i> Saussure | 7 ex. |
| <i>Hemipepsis glabrata</i> Van der Linden | 8 ex. |
| <i>Hemipepsis heros</i> Guérin, 1848 | 3 ex. |
| <i>Hemipepsis imperialis</i> Smith | 1 ex. |
| <i>Hemipepsis insulana</i> Wahis | 1 ex. |
| <i>Hemipepsis iodoptera</i> Stal | 11 ex. |
| <i>Hemipepsis sauteri</i> Wahis | 1 ex. |
| <i>Hemipepsis</i> sp. | 2 ex. |
| <i>Hemipepsis tamisieri</i> Guérin, 1848 | 6 ex. |
| <i>Hemipepsis ustulata ochroptera</i> Stal, 1857 | 2 ex. |
| <i>Hemipepsis variabilis</i> Arnold | 2 ex. |
| <i>Hemipepsis vespertilio</i> Gerstaecker | 5 ex. |
| <i>Hemipepsis vindex</i> Smith, 1855 | 1 ex. |
| <i>Homonotus balcanicus</i> Haupt, 1927 | 3 ex. |
| <i>Homonotus formosanus</i> Yasumatsu | 3 ex. |
| <i>Homonotus ruficaudis</i> Cameron | 1 ex. |
| <i>Homonotus sanguinolentus</i> Fabricius, 1793 | 20 ex. |
| <i>Irenangelus intrusus</i> Banks | 1 ex. |
| <i>Irenangelus pernix</i> Bingham | 1 ex. |
| <i>Java formosana</i> Wahis | 2 ex. |
| <i>Leptodialepis bipartita</i> Lepeletier | 1 ex. |
| <i>Leptodialepis formosanus</i> Wahis | 6 ex. |
| <i>Leptodialepis nicevillii</i> Bingham, 1896 | 15 ex. |
| <i>Leptodialepis variabilis variabilis</i> Wahis | 1 ex. |
| <i>Macromeris violacea</i> Lepeletier | 1 ex. |
| <i>Microphadnus pumilus</i> Costa, 1882 | 2 ex. |
| <i>Mygnumia chinensis</i> Smith | 8 ex. |

| | |
|---|--------|
| <i>Mygnumia consanguinea</i> Saussure | 1 ex. |
| <i>Mygnumia contumax</i> Gerstaecker | 1 ex. |
| <i>Mygnumia curvinervis</i> Cameron | 2 ex. |
| <i>Mygnumia matangensis</i> Cameron | 1 ex. |
| <i>Nanoclavelia leucopterus</i> Dahlbom, 1845 | 7 ex. |
| <i>Neagenia asiatica</i> Suster, i.l. | 11 ex. |
| <i>Notiochaes philadelphicus</i> Lepeletier | 2 ex. |
| <i>Notocyphus dorsalis arizonicus</i> Townes | 2 ex. |
| <i>Paraclavelia crudelis</i> Smith | 1 ex. |
| <i>Paraclazolia rhodesiensis</i> Arnold | 1 ex. |
| <i>Paracyphononyx alienus</i> Smith | 20 ex. |
| <i>Paracyphononyx beckeri</i> Cameron | 2 ex. |
| <i>Paracyphononyx difficilis</i> Bischoff | 7 ex. |
| <i>Paracyphononyx funereus</i> Lepeletier, 1845 | 2 ex. |
| <i>Paracyphononyx languidus</i> H..? | 1 ex. |
| <i>Paracyphononyx nutemmensis</i> Mayer | 1 ex. |
| <i>Paracyphononyx pedestris</i> Smith, 1855 | 6 ex. |
| <i>Paracyphononyx ruficrus</i> Klug | 9 ex. |
| <i>Paracyphononyx serraticornis</i> Taschenberg | 1 ex. |
| <i>Paracyphononyx zonatus</i> Illiger | 9 ex. |
| <i>Paraferreola dimidiata</i> Dahlbom | 2 ex. |
| <i>Paraferreola formosanus</i> Babiy, i.l. | 1 ex. |
| <i>Paraferreola manicata</i> Pallas | 3 ex. |
| <i>Paraferreola melanostoma</i> Cameron | 1 ex. |
| <i>Pepsis amabilis</i> Mocsáry | 2 ex. |
| <i>Pepsis amyntas</i> Mocsáry | 1 ex. |
| <i>Pepsis asteria</i> Mocsáry | 3 ex. |
| <i>Pepsis atrovirens</i> Lucas | 1 ex. |
| <i>Pepsis auranticornis</i> Lucas | 1 ex. |
| <i>Pepsis aurozonata</i> Smith | 1 ex. |
| <i>Pepsis basifusca</i> Lucas | 1 ex. |
| <i>Pepsis bonariensis</i> Lepeletier | 1 ex. |
| <i>Pepsis brasiliensis</i> Bréthes | 1 ex. |
| <i>Pepsis brevicornis</i> Bréthes | 1 ex. |
| <i>Pepsis caridei</i> Bréthes | 1 ex. |
| <i>Pepsis charon</i> Mocsáry | 2 ex. |
| <i>Pepsis chlorana</i> Mocsáry | 1 ex. |
| <i>Pepsis chrysothemis</i> Lucas, 1895 | 4 ex. |
| <i>Pepsis cinnabarina</i> Lucas | 2 ex. |
| <i>Pepsis completa</i> Smith | 2 ex. |
| <i>Pepsis culta</i> Bréthes | 1 ex. |
| <i>Pepsis cylindrica</i> Lucas | 1 ex. |
| <i>Pepsis defecta</i> Thunberg | 1 ex. |
| <i>Pepsis deuteroleuca</i> Smith | 2 ex. |
| <i>Pepsis diana</i> Mocsáry | 1 ex. |
| <i>Pepsis dimidiata</i> Fabricius, 1805 | 9 ex. |
| <i>Pepsis discolor</i> Taschenberg | 5 ex. |

| | |
|---|--------|
| <i>Pepsis domingensis</i> Lepeletier | 10 ex. |
| <i>Pepsis elevata</i> Fabricius | 2 ex. |
| <i>Pepsis erdmanni</i> Lucas | 1 ex. |
| <i>Pepsis erythroptera</i> Mocsáry | 1 ex. |
| <i>Pepsis eximia</i> Fabricius | 1 ex. |
| <i>Pepsis formosa</i> Say | 1 ex. |
| <i>Pepsis frivaldskii</i> Mocsáry | 1 ex. |
| <i>Pepsis fulgidipennis</i> Mocsáry | 7 ex. |
| <i>Pepsis glabripennis</i> Lucas | 1 ex. |
| <i>Pepsis grossa</i> Fabricius, 1789 | 8 ex. |
| <i>Pepsis heros</i> Fabricius | 7 ex. |
| <i>Pepsis ianthina</i> Erichson | 2 ex. |
| <i>Pepsis karschi</i> Lucas | 1 ex. |
| <i>Pepsis lampas</i> Lucas | 2 ex. |
| <i>Pepsis lepida</i> Mocsáry | 3 ex. |
| <i>Pepsis limbata</i> Guérin | 47 ex. |
| <i>Pepsis luteicornis</i> Fabricius | 1 ex. |
| <i>Pepsis lynchii</i> Bréthes | 1 ex. |
| <i>Pepsis marginata</i> Palisot de Beauvois, 1809 | 2 ex. |
| <i>Pepsis nessus</i> Lucas, 1895 | 1 ex. |
| <i>Pepsis nigrescens</i> Smith | 2 ex. |
| <i>Pepsis niphe</i> Mocsáry | 1 ex. |
| <i>Pepsis nupta</i> Lucas | 1 ex. |
| <i>Pepsis obliquerrugosa</i> Lucas | 3 ex. |
| <i>Pepsis optima</i> Smith | 2 ex. |
| <i>Pepsis patagonica</i> Bréthes | 1 ex. |
| <i>Pepsis planifrons</i> Lucas | 2 ex. |
| <i>Pepsis plutus</i> Erichson | 1 ex. |
| <i>Pepsis pretiosa</i> Dahlbom | 1 ex. |
| <i>Pepsis pulchripennis</i> Mocsáry | 7 ex. |
| <i>Pepsis reaumuri</i> Dahlbom | 9 ex. |
| <i>Pepsis rubra</i> Drury | 5 ex. |
| <i>Pepsis ruficornis</i> Fabricius | 10 ex. |
| <i>Pepsis sagana</i> Mocsáry | 8 ex. |
| <i>Pepsis sanguigutta</i> Christ | 2 ex. |
| <i>Pepsis schrottkyi</i> Bréthes | 1 ex. |
| <i>Pepsis seladonica</i> Dahlbom | 2 ex. |
| <i>Pepsis selene</i> Lucas | 1 ex. |
| <i>Pepsis sinnis</i> Lucas | 1 ex. |
| <i>Pepsis smaragdina</i> Dahlbom | 5 ex. |
| <i>Pepsis smaragdinula</i> Lucas | 6 ex. |
| <i>Pepsis</i> sp. | 6 ex. |
| <i>Pepsis sphinx</i> Lucas | 1 ex. |
| <i>Pepsis vaualba</i> Smith | 1 ex. |
| <i>Pepsis venusta</i> Smith, 1855 | 6 ex. |
| <i>Pepsis viridisetosa</i> Spinola | 1 ex. |
| <i>Pepsis xanthocera</i> Dahlbom | 3 ex. |

| | |
|---|---------|
| <i>Pepsis zelotarum</i> Lucas | 1 ex. |
| <i>Phanochilus ornatus</i> Banks | 1 ex. |
| <i>Platyderes fuscipennis</i> Babiý, i.l. | 1 ex. |
| <i>Platydialepsis fluctuata formosana</i> Wahis | 1 ex. |
| <i>Poecilagenia bicolorata</i> Susterá, i.l. | 2 ex. |
| <i>Poecilagenia formosana</i> Susterá, i.l. | 3 ex. |
| <i>Poecilagenia minor</i> Susterá, i.l. | 1 ex. |
| <i>Poecilagenia nigra</i> Susterá, i.l. | 4 ex. |
| <i>Poecilagenia rubricans</i> Lepeletier, 1845 | 1 ex. |
| <i>Poecilopompilus algidus fervidus</i> Smith | 2 ex. |
| <i>Poecilopompilus interruptus</i> Say, 1835 | 2 ex. |
| <i>Poecilopompilus interruptus cressoni</i> Banks | 2 ex. |
| <i>Poecilopompilus mixtus</i> Fabricius, 1798 | 4 ex. |
| <i>Pompilus apicatus</i> Provancher | 3 ex. |
| <i>Pompilus cinctellus</i> Van der Linden | 2 ex. |
| <i>Pompilus cinereus</i> Fabricius, 1775 | 287 ex. |
| <i>Pompilus exaltatus</i> Panzer | 2 ex. |
| <i>Pompilus expulsus</i> Schulz, 1906 | 2 ex. |
| <i>Pompilus formosanus</i> Babiý | 1 ex. |
| <i>Pompilus luctuosus</i> Cresson | 8 ex. |
| <i>Pompilus neglectus</i> Wesmael | 1 ex. |
| <i>Pompilus niger</i> Van der Linden | 3 ex. |
| <i>Pompilus obtusiventris</i> Dahlbom | 1 ex. |
| <i>Pompilus pectinipes</i> Van der Linden | 6 ex. |
| <i>Pompilus pusillus</i> Dahlbom | 4 ex. |
| <i>Pompilus scelestus</i> Cresson | 6 ex. |
| <i>Pompilus sericeus</i> Schiödte | 4 ex. |
| <i>Pompilus spinolae</i> Kohl | 1 ex. |
| <i>Priochilus captivus</i> Fabricius | 4 ex. |
| <i>Priochilus gracillimus</i> Smith | 2 ex. |
| <i>Priochilus pectoralis</i> Smith | 5 ex. |
| <i>Priochilus sericeifrons</i> Fox | 1 ex. |
| <i>Priochilus splendidulus</i> Fabricius | 1 ex. |
| <i>Priochilus veraepacis</i> Cameron | 1 ex. |
| <i>Priocnemella</i> sp. | 1 ex. |
| <i>Priocnemis agilis</i> Shuckard, 1837 | 78 ex. |
| <i>Priocnemis auripennis</i> Fabricius | 2 ex. |
| <i>Priocnemis baltica</i> Blüthgen, 1944 | 1 ex. |
| <i>Priocnemis bellieri</i> Sichel, 1860 | 7 ex. |
| <i>Priocnemis bifasciatus</i> Pérez | 1 ex. |
| <i>Priocnemis brunneus</i> Klug, 1834 | 1 ex. |
| <i>Priocnemis coeruleus</i> Taschenberg | 2 ex. |
| <i>Priocnemis consimilis</i> Costa | 7 ex. |
| <i>Priocnemis cordivalvata</i> Haupt, 1927 | 142 ex. |
| <i>Priocnemis coriacea</i> Dahlbom, 1843 | 79 ex. |
| <i>Priocnemis cotesi</i> Cameron | 1 ex. |
| <i>Priocnemis enslini</i> Haupt, 1927 | 3 ex. |

| | |
|---|---------|
| <i>Priocnemis exaltata</i> Fabricius, 1775 | 149 ex. |
| <i>Priocnemis faillai</i> Destefani, 1886 | 2 ex. |
| <i>Priocnemis fallax</i> Verhoeff, 1922 | 1 ex. |
| <i>Priocnemis fastigiata</i> Haupt, 1935 | 1 ex. |
| <i>Priocnemis fennica</i> Haupt, 1927 | 67 ex. |
| <i>Priocnemis flavipes</i> Guérin | 1 ex. |
| <i>Priocnemis germana</i> Cresson, 1867 | 2 ex. |
| <i>Priocnemis gracilis</i> Haupt, 1927 | 8 ex. |
| <i>Priocnemis gussakovskii</i> Lelej | 3 ex. |
| <i>Priocnemis hankoi</i> Moczar, 1944 | 11 ex. |
| <i>Priocnemis haupti</i> Moczar, 1955 | 1 ex. |
| <i>Priocnemis hirticeps</i> Guérin | 6 ex. |
| <i>Priocnemis hyalinata</i> Fabricius, 1793 | 159 ex. |
| <i>Priocnemis japonica</i> Gussakovskij | 60 ex. |
| <i>Priocnemis melanosoma</i> Kohl, 1880 | 2 ex. |
| <i>Priocnemis minuta</i> Van der Linden, 1827 | 136 ex. |
| <i>Priocnemis minutalis</i> Wahis, 1979 | 14 ex. |
| <i>Priocnemis oregona</i> Banks, 1933 | 5 ex. |
| <i>Priocnemis parvula</i> Dahlbom, 1845 | 19 ex. |
| <i>Priocnemis perturbator</i> Harris, 1780 | 370 ex. |
| <i>Priocnemis pogonioides</i> ichnusa Wolf | 23 ex. |
| <i>Priocnemis pogonioides pogonioides</i> Costa, 1883 | 1 ex. |
| <i>Priocnemis propinqua</i> Lepeletier, 1845 | 4 ex. |
| <i>Priocnemis pseudojaponica</i> Lelej | 2 ex. |
| <i>Priocnemis pusilla</i> Schiöde, 1837 | 91 ex. |
| <i>Priocnemis rothneyi</i> Cameron | 1 ex. |
| <i>Priocnemis rugosa</i> Susterai, 1922 | 3 ex. |
| <i>Priocnemis schioedtei</i> Haupt, 1927 | 38 ex. |
| <i>Priocnemis</i> sp. | 3 ex. |
| <i>Priocnemis sulci</i> Balthasar, 1943 | 1 ex. |
| <i>Priocnemis susterai</i> Haupt, 1927 | 11 ex. |
| <i>Priocnemis thaumastarius</i> Kohl | 4 ex. |
| <i>Priocnemis variegatus</i> Fabricius | 7 ex. |
| <i>Priocnemis vulgaris</i> Dufour, 1841 | 55 ex. |
| <i>Priocnessus fluctuatus</i> Cameron | 2 ex. |
| <i>Priocnessus nebulosus</i> Dahlbom, 1843 | 1 ex. |
| <i>Priocnessus obscurus</i> Wahis | 2 ex. |
| <i>Pseudopompilus humboldti</i> Dahlbom | 1 ex. |
| <i>Psorthaspis planata</i> Fox, 1892 | 2 ex. |
| <i>Salius odin</i> Strand, 1914 | 1 ex. |
| <i>Schistonyx umbrosus</i> Klug, 1834 | 7 ex. |
| <i>Sericopompilus neotropicalis</i> Cameron, 1893 | 2 ex. |
| <i>Sphictostethus gravesii</i> Haliday, 1837 | 3 ex. |
| <i>Spuridiophorus maculipennis</i> Arnold | 1 ex. |
| <i>Tachyagetes aemulans tenerifensis</i> Wolf, 1975 | 1 ex. |
| <i>Tachyagetes filicornis</i> Tournier, 1889 | 4 ex. |
| <i>Tachyagetes testaceoides</i> Wolf, 1982 | 3 ex. |

| | |
|---|-------|
| Tachypompilus analis Fabricius, 1781 | 2 ex. |
| Tachypompilus ferrugineus ferrugineus Say, 1824 | 1 ex. |
| Tachypompilus latus Smith | 1 ex. |
| Tachypompilus mendozae Dalla Torre | 1 ex. |
| Tachypompilus unicolor unicolor Banks, 1919 | 2 ex. |
| Teinotrachelus longicollis Bischoff | 1 ex. |
| Teinotrachelus ruficolor Arnold | 1 ex. |
| Telostegus inermis Brullé, 1832 | 3 ex. |
| Trachyglyptus spinosipes Turner | 1 ex. |