

The study of harmful benthic dinoflagellates started in the late 1970s when it was suspected that a benthic species, later named *Gambierdiscus toxicus*, was responsible for ciguatera fish poisoning, a type of human poisoning linked to the consumption of certain species of tropical reef fish. As the number of ciguatera fish poisoning incidents increases, and the distribution of toxin producing benthic taxa seems to be expanding, detailed understanding of the species diversity and the ability to accurately identify them is becoming increasingly important (chapter VII).

Dinoflagellate classification is currently undergoing changes and far from being settled, as new species and genera are discovered and systematic entities are rearranged. Many benthic dinoflagellate genera have unusual morphologies and appear to be only remotely related to known planktonic taxa, so that molecular phylogenetic analyses frequently show little statistical support for any relationship (chapter IV). Benthic species display unique thecal plate arrangements compared to planktonic species, e.g. *Adenoides*, *Amphidiniella*, *Cabra*, *Planodinium*, *Sabulodinium*, *Rhinodinium* (chapter III). Therefore, no classification on higher rank levels (e.g. family, order) was used throughout this book. Genera (and species within a genus) are presented in alphabetical order.

This book presents the first comprehensive identification help for benthic dinoflagellates. At the same time it

aims to lend support in order to improve monitoring efforts worldwide. About 190 species in 45 genera are presented in detail, illustrated with more than 200 color images, approximately 150 scanning electron micrographs, and more than 250 drawings.

This book is the first summary of our knowledge of benthic dinoflagellate species.

Dinoflagellates are important primary producers and symbionts, but, at the same time, also consumers and parasites. Species compositions of benthic habitats are quite distinct from those of planktonic habitats. Less than 10% of the approximately 2000 described extant dinoflagellate species appear to be benthic. They occur in different types of habitats (chapter II) and their morphology, their behavior, and some of their life cycles (chapter VI) seem to be well adapted to the benthic life-style. Information on their geographic distribution is still very limited and is compiled herein (chapter V).




Table of Contents

Greetings	4	<i>Herdmania</i>	109	'Dinotoms' – <i>Dinotrix</i> , <i>Durinskia</i> ,	
Foreword	5	<i>Heterocapsa</i>	111	<i>Galeidinium</i> , ' <i>Gymnodinium</i> ' <i>quadri-</i>	
Contents	7	<i>Katodinium</i>	112	<i>batum</i> , ' <i>Peridinium</i> ' <i>quinquecorne</i>	198
Acknowledgements	10	<i>Moestrupia</i>	115	Dinoflagellate taxa with cryptophyte-	
		<i>Ostreopsis</i>	116	(klepto)chloroplasts	198
I. Introduction	12	' <i>Peridinium</i> ' partim = new genus	126	The phytodinialean dinoflagellates	
		<i>Pileidinium</i>	128	('Phytodiniales')	198
II. 'Materials & Methods'	16	<i>Plagiodinium</i>	129		
Habitats	16	<i>Planodinium</i>	130	V. Biogeography	209
Sampling	16	<i>Polykrikos</i>	132		
Extraction = separation from		<i>Prorocentrum</i>	134	VI. Ecology	212
the substrate	19	<i>Pseudothecadinium</i>	152	Attachment	213
Fixation and Electron		<i>Pyramidodinium</i>	154	Life cycles	213
Microscopy (EM)	19	<i>Rhinodinium</i>	155	Tide pools	213
Culturing	21	<i>Roscoffia</i>	156	Vertical migration	214
Quantification	21	<i>Sabulodinium</i>	160	Blooms	214
		<i>Scrippsiella</i>	163	Spatial distribution	216
III. Taxonomy	22	<i>Sinophysis</i>	165	Temporal distribution	216
<i>Adenoides</i>	22	<i>Spiniferodinium</i>	173	Quantitative Data	217
<i>Alexandrium</i>	24	<i>Stylodinium</i>	175		
<i>Amphidiniella</i>	25	<i>Symbiodinium</i> spp.	177	VII. Toxins of benthic dinoflagellates and	
<i>Amphidiniopsis</i>	27	<i>Testudodinium</i>	178	benthic harmful algal blooms	218
<i>Amphidinium</i>	41	<i>Thecadinium</i>	180	Introduction	218
<i>Ankistrodinium</i>	56	<i>Togula</i>	188	<i>Gambierdiscus</i>	219
<i>Apicoporus</i>	58	<i>Vulcanodinium</i>	191	<i>Ostreopsis</i>	223
<i>Biecheleria</i>	61		193	<i>Coolia</i>	223
<i>Bispinodinium</i>	62	IV. Phylogeny and systematics	193	<i>Prorocentrum</i>	226
<i>Bysmatrum</i>	64	Phylogeny of the morphological	194	<i>Amphidinium</i>	227
<i>Cabra</i>	70	adaptations	194	<i>Alexandrium</i>	227
<i>Coolia</i>	74	<i>Amphidinium</i>	195	<i>Vulcanodinium</i>	227
<i>Dinotrix</i>	80	<i>Amphidiniopsis</i> , <i>Archaeoperidinium</i> ,	195		
<i>Durinskia</i>	82	<i>Herdmania</i> – Peridiniales	195	References	234
<i>Galeidinium</i>	85	<i>Cabra</i> , <i>Rhinodinium</i> , <i>Roscoffia</i> –	196	Taxonomic index	266
<i>Gambierdiscus</i>	86	<i>Podolampadaceae</i>	196	Useful web pages	272
<i>Glenodinium</i>	95	<i>Coolia</i> , <i>Gambierdiscus</i> , <i>Ostreopsis</i> –	196	Picture credits	273
<i>Gymnodinium</i>	96	<i>Gonyaulacales</i>	196	Authors' Addresses	274
<i>Gyrodinium</i>	103	<i>Prorocentrum</i> & <i>Adenoides</i>	197		
<i>Halostylodinium</i>	107	<i>Sinophysis</i> & <i>Sabulodinium</i>	197		

Order Form

I/we would like to order from

E. Schweizerbart'sche Verlagsbuchhandlung (Nägele u. Obermiller)
 Johannesstr. 3A | 70176 Stuttgart | Germany
 Phone +49 (0)711 351456 0 | FAX +49 (0)711 351456 99
 order@schweizerbart.de | www.schweizerbart.de/9783510614028

..... Copies

Kleine Senckenberg-Reihe 54
 Mona Hoppenrath | Shauna A. Murray | Nicolas Chomérat | Takeo Horiguchi
Marine benthic dinoflagellates – unveiling their worldwide biodiversity
 14.8 x 21 cm | paperback | 19.90 € | ISBN 978-3-510-61402-8

Name

Address

.....

Date

Signature

